

Standard Normal (z) Distribution

Showing area between 0 and z; that is,  $P(0 \leq x \leq z)$

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.0000	.0040	.0080	.0120	.0160	.0199	.0239	.0279	.0319	.0359
0.1	.0398	.0438	.0478	.0517	.0557	.0596	.0636	.0675	.0714	.0753
0.2	.0793	.0832	.0871	.0910	.0948	.0987	.1026	.1064	.1103	.1141
0.3	.1179	.1217	.1255	.1293	.1331	.1368	.1406	.1443	.1480	.1517
0.4	.1554	.1591	.1628	.1664	.1700	.1736	.1772	.1808	.1844	.1879
0.5	.1915	.1950	.1985	.2019	.2054	.2088	.2123	.2157	.2190	.2224
0.6	.2257	.2291	.2324	.2357	.2389	.2422	.2454	.2486	.2517	.2549
0.7	.2580	.2611	.2642	.2673	.2704	.2734	.2764	.2794	.2823	.2852
0.8	.2881	.2910	.2939	.2967	.2995	.3023	.3051	.3078	.3106	.3133
0.9	.3159	.3186	.3212	.3238	.3264	.3289	.3315	.3340	.3365	.3389
1.0	.3413	.3438	.3461	.3485	.3508	.3531	.3554	.3577	.3599	.3621
1.1	.3643	.3665	.3686	.3708	.3729	.3749	.3770	.3790	.3810	.3830
1.2	.3849	.3869	.3888	.3907	.3925	.3944	.3962	.3980	.3997	.4015
1.3	.4032	.4049	.4066	.4082	.4099	.4115	.4131	.4147	.4162	.4177
1.4	.4192	.4207	.4222	.4236	.4251	.4265	.4279	.4292	.4306	.4319
1.5	.4332	.4345	.4357	.4370	.4382	.4394	.4406	.4418	.4429	.4441
1.6	.4452	.4463	.4474	.4484	.4495	.4505	.4515	.4525	.4535	.4545
1.7	.4554	.4564	.4573	.4582	.4591	.4599	.4608	.4616	.4625	.4633
1.8	.4641	.4649	.4656	.4664	.4671	.4678	.4686	.4693	.4699	.4706
1.9	.4713	.4719	.4726	.4732	.4738	.4744	.4750	.4756	.4761	.4767
2.0	.4772	.4778	.4783	.4788	.4793	.4798	.4803	.4808	.4812	.4817
2.1	.4821	.4826	.4830	.4834	.4838	.4842	.4846	.4850	.4854	.4857
2.2	.4861	.4864	.4868	.4871	.4875	.4878	.4881	.4884	.4887	.4890
2.3	.4893	.4896	.4898	.4901	.4904	.4906	.4909	.4911	.4913	.4916
2.4	.4918	.4920	.4922	.4925	.4927	.4929	.4931	.4932	.4934	.4936
2.5	.4938	.4940	.4941	.4943	.4945	.4946	.4948	.4949	.4951	.4952
2.6	.4953	.4955	.4956	.4957	.4959	.4960	.4961	.4962	.4963	.4964
2.7	.4965	.4966	.4967	.4968	.4969	.4970	.4971	.4972	.4973	.4974
2.8	.4974	.4975	.4976	.4977	.4977	.4978	.4979	.4979	.4980	.4981
2.9	.4981	.4982	.4982	.4983	.4984	.4984	.4985	.4985	.4986	.4986
3.0	.4987	.4987	.4987	.4988	.4988	.4989	.4989	.4989	.4990	.4990

T.O.H.	$\alpha = .10$	$\alpha = .05$	$\alpha = .01$	C.I.	z
One-tailed	1.282	1.645	2.327	90%	1.645
Two-tailed	1.645	1.960	2.575	95%	1.960
				99%	2.575

Critical Values of the t Distribution

Showing CV given df and  $\alpha$ ; that is,  $P(t \geq CV) = \alpha$

df	$\alpha$				
	.005 (1 tail)	.01 (1 tail)	.025 (1 tail)	.05 (1 tail)	.10 (1 tail)
	.01 (2 tails)	.02 (2 tails)	.05 (2 tails)	.10 (2 tails)	.20 (2 tails)
1	63.657	31.821	12.706	6.314	3.078
2	9.925	6.965	4.303	2.920	1.886
3	5.841	4.541	3.182	2.353	1.638
4	4.604	3.747	2.776	2.132	1.533
5	4.032	3.365	2.571	2.015	1.476
6	3.707	3.143	2.447	1.943	1.440
7	3.500	2.998	2.365	1.895	1.415
8	3.355	2.896	2.306	1.860	1.397
9	3.250	2.821	2.262	1.833	1.383
10	3.169	2.764	2.228	1.812	1.372
11	3.106	2.718	2.201	1.796	1.363
12	3.054	2.681	2.179	1.782	1.356
13	3.012	2.650	2.160	1.771	1.350
14	2.977	2.625	2.145	1.761	1.345
15	2.947	2.602	2.132	1.753	1.341
16	2.921	2.584	2.120	1.746	1.337
17	2.898	2.567	2.110	1.740	1.333
18	2.878	2.552	2.101	1.734	1.330
19	2.861	2.540	2.093	1.729	1.328
20	2.845	2.528	2.086	1.725	1.325
21	2.831	2.518	2.080	1.721	1.323
22	2.819	2.508	2.074	1.717	1.321
23	2.807	2.500	2.069	1.714	1.320
24	2.797	2.492	2.064	1.711	1.318
25	2.787	2.485	2.060	1.708	1.316
26	2.779	2.479	2.056	1.706	1.315
27	2.771	2.473	2.052	1.703	1.314
28	2.763	2.467	2.048	1.701	1.313
29	2.756	2.462	2.045	1.699	1.311
z	2.575	2.327	1.960	1.645	1.282

Critical Values of the Pearson Correlation Coefficient r

n	$\alpha = .05$	$\alpha = .01$	Least Squares Regression
4	.950	.999	
5	.878	.959	$\bar{X} = \Sigma X/n$ $\bar{Y} = \Sigma Y/n$
6	.811	.917	
7	.754	.875	
8	.707	.834	$SS_{XX} = \Sigma (X^2) - (\Sigma X)^2/n$
9	.666	.798	
10	.632	.765	$SS_{YY} = \Sigma (Y^2) - (\Sigma Y)^2/n$
11	.602	.735	
12	.576	.708	$SS_{XY} = \Sigma (XY) - (\Sigma X)(\Sigma Y)/n$
13	.553	.684	
14	.532	.661	
15	.514	.641	$\beta_1' = SS_{XY} / SS_{XX}$ (slope)
16	.497	.623	
17	.482	.606	$\beta_0' = \bar{Y} - \beta_1' \bar{X}$ (intercept)
18	.468	.590	
19	.456	.575	
20	.444	.561	$Y' = \beta_0' + \beta_1' X$
25	.396	.505	
30	.361	.463	$r = SS_{XY} / \sqrt{SS_{XX}SS_{YY}}$
35	.335	.430	
40	.312	.402	
45	.294	.378	
50	.279	.361	
60	.254	.330	
70	.236	.305	
80	.220	.286	
90	.207	.269	
100	.196	.256	

To test  $H_0: \rho = 0$  against  $H_1: \rho \neq 0$ ,  
 reject  $H_0$  if  $|r| > CV$ . Note: if we can  
 reject  $H_0$  then the model is useable.

Chi-square ( $\chi^2$ ) Distribution

Test for Goodness-of-Fit

df	Critical value (CV)	
	= .05	= .01
1	3.841	6.635
2	5.991	9.210
3	7.815	11.345
4	9.488	13.277
5	11.071	15.086
6	12.592	16.812
7	14.067	18.475
8	15.507	20.090
9	16.919	21.666
10	18.307	23.209
11	19.675	24.725
12	21.026	26.217
13	22.362	27.688
14	23.685	29.141
15	24.996	30.578
16	26.296	32.000
17	27.587	33.409
18	28.869	34.805
19	30.144	36.191
20	31.410	37.566
21	32.671	38.932
22	33.924	40.289
23	35.172	41.638
24	36.415	42.980
25	37.652	44.314
26	38.885	45.642
27	40.113	46.963
28	41.337	48.278
29	42.557	49.588
30	43.773	50.892
40	55.758	63.691
50	67.505	76.154
60	79.082	88.379
70	90.531	100.425
80	101.879	112.329
90	113.145	124.116
100	124.342	135.807

$$\chi^2 = \sum \left[ \frac{(O - E)^2}{E} \right]$$

Use df = clms-1

Reject Ho: fit as stated

if  $\chi^2 > CV$

Test for Independence

$$E = \frac{(\text{row total}) * (\text{clm total})}{(\text{grand total})}$$

Use df = (rows-1) \* (clms-1)

Reject Ho: independent

if  $\chi^2 > CV$

Showing  $P(\chi^2 \geq CV) = \alpha$